

intercalators studied) this difference is explained by the importance of the charge transfer term which is not included in the AMBER potential. The Hartree-Fock and DFT/B3LYP methods not covering the dispersion energy fail completely to describe any energy minimum at the potential energy curve of the E...AT complex and these methods thus cannot be recommended for a study of intercalation process. On the other hand, a modified version of DFT method which covers London dispersion energy yields for all complexes very good stabilization energies well comparable with referenced *ab initio* data. Besides vertical dependence of interaction energy twist dependence of interaction energy was also investigated by both, reference correlated *ab initio* method as well as empirical potentials. It is concluded that despite the charged (E +1, D +1, DAPI +2) or polar (EL) character of intercalators investigated it is the dispersion energy which predominantly contributes to the stability of intercalator...DNA base pair complexes. Any procedure which does not cover dispersion energy is thus not suitable for studying the process of intercalation.

1. D. Řeha, M. Kabeláč, F. Ryjáček, J. Šponer, J. E. Šponer, M. Elstner, S. Suhai, P. Hobza, *J. Am. Chem. Soc.*, **124** (2002) 3366-3376.

ANALYSIS OF INTERACTIONS IN COMPLEXES OF HIV-1 PROTEASE AND ITS PEPTIDOMIMETIC INHIBITOR

T. Skálová, H. Petroková, J. Hašek, J. Dohnálek, E. Buchtelová, J. Dušková

Institute of Macromolecular Chemistry, Academy of Sciences of the Czech Republic, Heyrovského nám. 2, 162 06 Praha 6, Czech Republic

HIV-1 protease is a 22 kDa protein of the human immunodeficiency virus. The function of this protein is to cleave polyprotein of immature virus and thus to contribute to formation of active matured virus. Inhibition of the protease is therefore one of possible ways of fighting with disease AIDS, caused by the human immunodeficiency virus.

Our research was focused on interaction analysis of HIV-1 protease and its peptidomimetic inhibitor Boc-Phe- [CH₂CH₂NH]-Phe-Glu-Phe-NH₂, denoted as OE. The inhibitor was developed in the laboratory of J. Konvalinka (Institute of Organic Chemistry and Biochemistry, Academy of Sciences CR). Native and mutant (A71V, V82T, I84V) HIV-1 protease were expressed and purified in the laboratories of J. Sedláček (Institute of Molecular Genetics, Academy of Sciences CR) and J. Konvalinka. In our research group, crystallization of complexes of OE with native and mutant protease was performed, X-ray diffraction of crystals on the synchrotron source of radiation was measured and structures of both complexes were determined ([1], [2]).

As a result, we have two structures with *R*-factors 18 % (native protease complex, diffraction limit 2.45 Å) and 20.3 % (mutant protease complex, diffraction limit 2.2 Å). Both complexes crystallized in space group P61 and in-

hibitor OE was found in the active site in two approximately C₂ symmetrical positions, following thus pseudo-symmetry of the protease. This fact makes interpretation of interactions between the protease and inhibitor more difficult. Therefore, standard structural analysis of contacts between the protease and inhibitor was completed by two energy analyses of interactions in the active site. The inhibitor binding modes to both proteases are similar from the structural point of view and interpretation of small details could be ambiguous. However, energy analysis of both complexes confirms the interpretation of changes caused by mutation of the protease. Mutated residue Thr 182 forms an aromatic hydrogen bond to the inhibitor phenyl group in P1 position. Mutation I84V causes a decrease in van der Waals interaction between residue 84 and the OE inhibitor.

The research was supported by the Grant Agency of the Academy of Sciences of the Czech Republic (projects A4050811/1998 and B4050312/2003) and by the Academy of Sciences of the Czech Republic (project AVOZ4050913).

1. T. Skálová, J. Hašek, J. Dohnálek, H. Petroková, E. Buchtelová, Mutant HIV-1 protease complexed with tetrapeptide inhibitor. Preliminary report, *Acta Phys. Pol. A*, **101** (2002), 659-663.
2. H. Petroková, unpublished results.

MEMBRANE PSEUDO-CRYSTAL STRUCTURES IN PSSU-IPT TOBACCO CHLOROPLASTS

Helena Synková¹, Renáta Pechová³, Michal Hušák², František Vácha², Pavel Šiffel²

¹*Institute of Experimental Botany, Academy of Sciences of the Czech Republic, Na Karlovce 1a, CZ-160 00 Praha 6*

²*Institute of Physical Biology, University of South Bohemia, Branišovská 31, 370 05 České Budějovice*

³*Department of Plant Physiology, Faculty of Sciences, Charles University, Viničná 5, CZ-128 44 Praha 2, Czech Republic*

Our study is focused on native pseudo-crystalline structures, which were observed in chloroplasts of transgenic tobacco overproducing plant hormones cytokinins. The structures were not positively identified until now. We suppose that they are formed by light harvesting protein (LHC) aggregating in a form of 2D crystal, which then constitute membrane stacks. Our hypothesis is supported by fluorescence emission spectra, which showed certain bands corresponding to LHC aggregates and higher emission of chlorophyll *b* in chloroplasts isolated from transgenic plants.

The aim of this experiment was the estimation of relative size of pseudo-crystals compared to chloroplast and the size of basic cell unit, which can be determined from analysis of TEM images from ultrathin sections of leaves and isolated chloroplast suspensions.

Transgenic tobacco containing a supplementary iptene under a control of the promoter for the small subunit of RuBPCO (*Pssu-ipt*) was grown as grafts on non-transgenic