

evaluating the dynamics and energetics of molecular models of RNA motifs constructed by phylogenetic analysis and isostericity principles.

1. C. C. Correll, B. Freeborn & P. B. Moore T. A. Steitz. *Cell*, **91** (1997) 705-712.
2. M. Lu & T. A. Steitz. *Proc. Natl. Acad. Sci. U S A*, **97** (2000) 2023-2028.
3. N. B. Leontis & E. Westhof. *RNA*, **4** (1998) 1134-1153.

THE EFFECT OF WATER SOLVENT ON THEORETICAL NMR SPIN - SPIN COUPLINGS IN DNA: IMPROVEMENT OF CALCULATED VALUES BY APPLICATION OF TWO SOLVENT MODELS

V. Sychrovský¹, B. Schneider¹, P. Hobza¹, L. Židek², V. Sklenář²

¹*J. Heyrovský Institute of Physical Chemistry, Academy of Sciences of the Czech Republic, and Center for Complex Molecular Systems and Biomolecules, Dolejškova 3, 18223, Praha 8, Czech Republic*

²*Laboratory of Biomolecular Structure and Dynamics and National Centre for Biomolecular Research, Faculty of Science, Masaryk University, Kotlářská 2, CZ-611 37, Brno, Czech Republic*

The NMR indirect spin - spin coupling constants calculated in DNA base pairs are compared with one-bond $^1J(X,H)$, $^1J(C,X)$ and two-bond $2J(X,H)$, ($X = C, N$) coupling constants measured in DNA hairpin molecule d(GCGAAGC) [1]. The two theoretical models of solvent: explicit and Polarized Continuum, were introduced. Both explicit and PCM solvent model provide a similar improvement of the calculated J constants when compared to the experimental data. The mean absolute deviation between the calculated and experimental couplings is significantly reduced by solvent inclusion, from 1.7 to 1.1 Hz for guanine, from 2.4 to 0.6 Hz for cytosine, and from 2.3 to 1.6 Hz for adenine. The strongest solvent shift was calculated for the $^1J(C,H)$ coupling constants, particularly for the (C8,H8) coupling in guanine (6.1 Hz) and adenine, and the (C5,H5) and (C6,H6) couplings in cytosine. These changes in $^1J(C,H)$ coupling seem to correspond to the charge transfer from water bonding and lone pair orbitals to the guanine anti-bonding C8-H8, C8-N7, and C8-N9 orbitals. From the close agreement between the calculated and experimental coupling constants we can deduce that the C-H group of the hairpin bases is extensively hydrated even when interaction with solvent is rather weak and non - specific.

1. V. Sychrovský, B. Schneider, P. Hobza, L. Židek, V. Sklenář, *Physical Chemistry Chemical Physics*, in press.

LECTURES - MARCH 15

STRUCTURAL BASIS OF HIV-1 AND HIV-2 PROTEASE INHIBITION BY A MONOCLONAL ANTIBODY

P. Řezáčová¹, J. Lescar², J. Brynda¹, M. Fábry¹, M. Hořejší¹, G.A. Bentley³ and J. Sedláček¹

¹*Dep. of Gene Manipulation, Institute of Molecular Genetics, Academy of Sciences of the Czech Republic, Flemingovo nám. 2, Prague 6, Czech Republic*

²*School of Biological Sciences, Nanyang Walk, Blk 5, Level 3, Singapore 637616*

³*Unité d'Immunologie Structurale, Institut Pasteur, 25 rue du Dr. Roux, 75724 Paris, France*

Since the demonstration that the Human Immunodeficiency Virus protease (HIV PR) is essential in the viral life cycle [1], this enzyme has become one of the primary targets for antiviral drug design. With the objective of probing the structural stability of HIV PR and the eventual design of potential inhibitors directed to regions other than the active site, we have examined the effects of anti-HIV-1 PR monoclonal antibodies (mAbs) on the catalytic activity of the protease [2, 3]. Monoclonal antibody mAb1696, although raised against the HIV-1 PR, inhibits the catalytic activity of both the HIV-1 and HIV-2 enzymes with inhibition constants in nanomolar range [3] and cross-reacts with peptides comprising the N-terminus of the HIV protease (residues 1 to 7). The N-terminal region is essential for dimerization of monomers and thus forming the active HIV PR dimer.

To investigate further the mechanism of HIV PR inhibition by mAb1696, a recombinant single-chain Fv fragment (scFv) that contains heavy and light chain variable domains of mAb1696 joined by a flexible linker have been prepared and used for crystallographic studies [4]. Although the crystallization of scFv1696 complexed with the HIV proteases has not yet been successful, the formation of these complexes has been demonstrated in solution. As an alternative approach to studying the antigenic recognition by the mAb 1696 at the structural level, we have crystallized complexes of scFv1696 with the respective epitope peptide fragments of the HIV protease. The three-dimensional structure of the complex formed between scFv1696 and epitope peptide PQITLWQRR (corresponding to the N-terminus of HIV-1 PR) has been solved and refined at 2.70 Å resolution [4]. The structure of the complex formed between scFv1696 and epitope peptide PQFSLWKR (corresponding to the N-terminus of HIV-2 PR) has been solved and refined at 1.88 Å resolution.

Interactions of peptides with scFv1696 have been analyzed and compared in both structures. On the basis of the interactions seen in the complex, the cross-reactivity between mAb1696 and the HIV-1 and HIV-2 protease and their N-terminal peptides can be explained. The broad reactivity between mAb 1696 on one side and the HIV-1 PR and HIV-2 PR derives from the invariance or conservation